CERN-PH-TH-2016-128

The Higgs of the Higgs and the diphoton channel

Kristjan Kannike,[a] **Giulio Maria Pelaggi,**[b] **Alberto Salvio,**[c] **Alessandro Strumia**[a,b,c]

[a] *National Institute of Chemical Physics and Biophysics, Rävala 10, 10143 Tallinn, Estonia*

[b] *Dipartimento di Fisica dell'Università di Pisa and INFN, Italy*

[c] *CERN, Theory Division, Geneva, Switzerland*

arXiv:1605.08681v2 [hep-ph] 22 Jul 2016

Abstract

LHC results do not confirm conventional natural solutions to the Higgs mass hierarchy problem, motivating alternative interpretations where a hierarchically small weak scale is generated from a dimension-less quantum dynamics. We propose weakly and strongly-coupled models where the field that breaks classical scale invariance giving mass to itself and to the Higgs is identified with a possible new resonance within the LHC reach. As an example, we identify such resonance with the 750 GeV diphoton excess recently reported by ATLAS and CMS. Such models can be extrapolated up to the Planck scale, provide Dark Matter candidates and eliminate the SM vacuum instability.

Contents

1 Introduction **2**

2 Weakly coupled models **3**
2.1 Model of type g) 5
2.2 Model of type λ) 9

3 Strongly coupled models **13**
3.1 The diphoton as a TCη 13
3.2 The diphoton as a TC-dilaton or TCσ 15

4 Conclusions **16**

A RGE for model g) **17**

B RGE for model λ) **18**

1 Introduction

Conventional solutions to the hierarchy problem have been disfavoured by experimental bounds that relegated them to fine-tuned corners of their parameter space. This situation motivated a reconsideration of the hierarchy problem: no physical effects are associated with quadratically divergent corrections to the Higgs squared masses, so maybe we are over-interpreting quantum field theory when we require theories that tame such quadratic divergences.

The hierarchy problem is bypassed if there are no particles much heavier than the Higgs and significantly coupled to it, such that physical corrections to the Higgs mass are naturally small. This heretic context was dubbed 'finite naturalness' in [1].

Within the general finite naturalness scenario an interesting sub-set of theories are those described by dimension-less Lagrangians, that assume that massive parameters do not exist at fundamental level, such that all mass scales in nature are generated dynamically, like the QCD scale in the Standard Model. Strictly speaking, finite-naturalness does not necessarily require the absence of masses in the Lagrangian. What makes the dynamical generation of masses attractive is the fact that it can lead to a separation of scales, which depends exponentially on dimensionless couplings. Therefore, this specific setup allows us to justify why the weak scale is many orders of magnitude smaller than the Planck scale, which itself might be generated by dimension-less dynamics, with important implications for inflation [2].

Notice that a dimension-less Lagrangian corresponds to a classical scale invariant model. We do not wish to preserve scale invariance at the quantum level because we eventually have to generate the observed scales. When the couplings of the theory are small, classical scale invariance is also an approximate symmetry of the model, otherwise it is just the requirement that the masses are dynamically generated (with the motivations stated above).

Various models where the weak scale arises in this way have been proposed. They can be classified in two categories:

1. The weak scale is the scale where a scalar quartic coupling λ runs negative, inducing vacuum expectation values *à la* Coleman-Weinberg [3–5].

2. The weak scale is the scale where a gauge coupling g runs to non-perturbative values, inducing condensates [6, 7].

It is interesting to ask whether these theoretical frameworks could lead to something visible at the LHC. A particularly clean channel, which has therefore a great discovery potential, is the one with two photons in the final state. We therefore study models where one of the required new particles can be identified with a new resonance F decaying into two photons (diphoton). As an example, we treat in some detail the case in which such resonance is identified with the recent diphoton excess at 750 GeV reported by ATLAS and CMS [8]. Other numerical choices are of course possible and our work will remain valuable even if such excess will turn out to be a statistical fluctuation.

Connections with conventional solutions to the hierarchy problem have been explored by authors who tried to identify F with one or another supersymmetric particle [9, 10] or with resonances of composite Higgs scenarios [9, 11].

References [12, 13] tried to incorporate the diphoton hinted by ATLAS and CMS in the weakly-coupled framework; however, in both cases couplings are so large that the attempted models hit Landau poles just above the weak scale, such that no hierarchy is dynamically generated.[1]

In section 2 we present weakly-coupled dimensionless models where F is the field that dynamically generates the weak scale, while running down from the Planck scale. In section 3 we present strongly coupled dimension-less models, where, among other things, the diphoton excess is reproduced. In section 4 we present our conclusions.

2 Weakly coupled models

Various extensions of the SM where the weak scale is generated *à la* Coleman-Weinberg have been proposed in the literature. They can be divided into two main categories, depending on which correction renormalises a quartic coupling λ_F down to negative values at low energy, such that the dimension-less potential $\lambda_F F^4$ develops a minimum: either

λ) corrections due to other scalar quartics [4]; or

g) corrections due to a gauge coupling [3, 5].[2]

[1] Furthermore [12] also contain some explicit mass term. The model in [14] can be extrapolated up to infinite energy, but it employs explicit mass terms. It is possible that these mass terms could be generated at a scale much higher than the weak scale. But finding an example of this sort goes beyond the scope of the present paper.

[2] Yukawa couplings have the opposite effect of making a quartic larger at low energy.

name	spin	$\mathrm{SU}(3)_c$	$\mathrm{SU}(2)_L$	$\mathrm{U}(1)_Y$	$\mathrm{SU}(N)$
X	0	1	1	0	1
S	0	1	1	0	N
$\mathcal{U}$	1/2	$\bar{3}$	1	$-2/3$	N
$\mathcal{U}^c$	1/2	3	1	$+2/3$	$\bar{N}$

Table 1: *Beyond the Standard Model field content of the model of type g) of section 2.1.*

Models of type λ) are more problematic than models of type g), because a non-abelian gauge coupling g can be sizeable without implying nearby Landau poles, while a sizeable quartic coupling drives itself to larger values at higher energies. Models of type λ) in the literature mitigate this effect by having a large number N of smaller quartics, and we will follow this strategy. We present in section 2.1 (section 2.2) models of type g) (of type λ) where the Coleman-Weinberg field is the diphoton, and where the RGE can be extrapolated up to the Planck scale, such that a large hierarchy is dynamically generated.

Before starting, we mention a broader — but less interesting — class of scale-invariant models, where the diphoton is added as an ad-hoc extra field that does not play a key role in the dynamical generation of the weak scale. Roughly, one can choose any one of the N diphoton models proposed in the literature, and any one of the M models that dynamically generated the weak scale, and combine them into $N \times M$ models.

For example, there is no obstacle in combining the dimension-less model in [3] (an extra $\mathrm{SU}(2)$ gauge interaction with an extra scalar doublet S, that acquires a vacuum expectation value through the Coleman-Weinberg mechanism) with the 'diphoton everybody's model' [9] (the diphoton is an extra scalar singlet F coupled to extra charged scalars X or fermions ψ). The SU(2) massive vectors are DM candidates [3].

What are the generic features of this class of ad-hoc models? The new charged particles cannot be arbitrarily heavier than the Higgs mass M_h, otherwise they contribute to it unnaturally [1]:

$$\begin{array}{ll} M_\psi \lesssim 4\pi M_h/g_{1,2}^2 & \text{for fermions,} \\ M_X \lesssim M_h/g_{1,2}^2 & \text{for scalars,} \end{array} \qquad (1)$$

where $g_{1,2}$ are the electroweak gauge couplings. The measured diphoton rate indeed suggests new particles with sub-TeV mass [9]. Furthermore, in the context of finite naturalness, the existence of extra colored fermions below a few TeV is necessary if the QCD θ problem is solved by an KSVZ axion model [1].

We now propose more interesting — but more constrained — models where the diphoton F is the particle that dynamically acquires the vacuum expectation value that induces the electroweak scale.

2.1 Model of type g)

We extend the SM by adding an extra gauge group $\mathrm{SU}(N)$ and the extra fields listed in table 1: two scalars S and X, a quark $\mathcal{U}$ with the same quantum numbers of the SM right-handed up quarks that fill a N of $\mathrm{SU}(N)$, plus the conjugated fermion $\mathcal{U}^c$ in order to form a vector-like quark. The dimension-less Yukawa couplings are

$$\mathscr{L}_Y = \mathscr{L}_Y^{\rm SM} + (y_S S\mathcal{U}^c U + y_X\, X\mathcal{U}\mathcal{U}^c + \text{h.c.}), \tag{2}$$

where U is the right-handed up quark of the SM. The dimension-less potential of the theory is

$$V(H,S,X) = \lambda_H|H|^4 + \lambda_X X^4 + \lambda_S|S|^4 - \lambda_{HS}|H|^2|S|^2 - \lambda_{HX}|H|^2X^2 - \lambda_{SX}|S|^2X^2. \tag{3}$$

The tree-level potential is positive, $V \geq 0$, when the quartic couplings satisfy [15]

$$\begin{cases} \lambda_H \geq 0, \qquad \lambda_S \geq 0, \qquad \lambda_X \geq 0, \\ \bar\lambda_{HS} \equiv -\lambda_{HS} + 2\sqrt{\lambda_H\lambda_S} \geq 0, \\ \bar\lambda_{HX} \equiv -\lambda_{HX} + 2\sqrt{\lambda_H\lambda_X} \geq 0, \\ \bar\lambda_{SX} \equiv -\lambda_{SX} + 2\sqrt{\lambda_S\lambda_X} \geq 0, \\ -\sqrt{\lambda_H}\lambda_{XS} - \sqrt{\lambda_S}\lambda_{HX} - \sqrt{\lambda_X}\lambda_{HS} + 2\sqrt{\lambda_H\lambda_S\lambda_X} + \sqrt{\bar\lambda_{HS}\bar\lambda_{HX}\bar\lambda_{SX}} \geq 0. \end{cases} \tag{4}$$

The RGEs of the model are listed in appendix A. Notice that λ_{HS} is unavoidably generated from the Yukawa couplings, if the top is the quark mostly coupled to the new states $\mathcal{U}$.

Masses

When the field S dynamically acquires a vacuum expectation value breaking $\mathrm{SU}(N) \to \mathrm{SU}(N-1)$ (to nothing if $N = 2$ [3]), H and X too can acquire vacuum expectation values, in view of their λ_{HS} and λ_{SX} quartic couplings to S. In the unitary gauge there are three physical scalars:

$$S = (v_S + \frac{s}{\sqrt{2}}, 0, \ldots), \qquad X = v_X + x, \qquad H = (v + \frac{h}{\sqrt{2}}, 0). \tag{5}$$

We assume that λ_{HS} and λ_{HX} are negligible, and that scale invariance gets broken when the RGE running of λ_S, dominated by the gauge coupling g, violates the condition $\bar\lambda_{SX} \geq 0$ of eq. (3). We can approximate the one loop potential by inserting a running

$$\lambda_S(S) = \frac{\lambda_{SX}^2}{4\lambda_X} + \beta_{\lambda_S} \ln\frac{|S|}{S_*} \tag{6}$$

in the tree-level potential of eq. (3). Here S_* is the scale at which the stability condition $\bar\lambda_{SX} \geq 0$ is violated. For $|S|$ around S_* there is an approximately flat direction, which is lifted by quantum correction (the log term in eq. (6)) as illustrated in fig.1. This generates the following absolute minimum of the potential

$$v_S = \frac{S_*}{e^{1/4}}, \qquad v_X = v_S\sqrt{\frac{\lambda_{SX}}{2\lambda_X}}, \tag{7}$$



Figure 1: *Potential defined in eq. (3) for λ_S chosen as in eq. (6) and $H = 0$. We set $\beta_{\lambda_S} \simeq 0.12$, $\lambda_X \simeq 0.030$ and $\lambda_{SX} \simeq 0.071$.*

which is visible in fig.1. There is another absolute minimum with the sign of v_X switched, but we assume $v_X > 0$ without loss of generality. The scalar mass matrix at the minimum in the (x, s) basis is

$$2v_S^2 \begin{pmatrix} 2\lambda_{SX} & -\sqrt{\lambda_{SX}^3/\lambda_X} \\ -\sqrt{\lambda_{SX}^3/\lambda_X} & \beta_{\lambda_S} + \lambda_{SX}^2/2\lambda_X \end{pmatrix}. \tag{8}$$

Neglecting the small mixing with the Higgs, the mass eigenstates are the diphoton F and a similar heavier scalar F'

$$\begin{cases} F = s\cos\theta + x\sin\theta \\ F' = x\cos\theta - s\sin\theta \end{cases} \qquad \text{where} \qquad \tan 2\theta = \frac{4\lambda_{SX}^{3/2}\lambda_X^{1/2}}{4\lambda_X\lambda_{SX} - \lambda_{SX}^2 - 2\lambda_X\beta_{\lambda_S}}. \tag{9}$$

Their masses are

$$M_{F,F'}^2 = \frac{v_S^2}{2\lambda_X}\left[A \mp \sqrt{A^2 - 32\beta_{\lambda_S}\lambda_X^2\lambda_{SX}}\right], \qquad A \equiv 4\lambda_X\lambda_{SX} + \lambda_{SX}^2 + 2\lambda_X\beta_{\lambda_S}. \tag{10}$$

The Higgs mass is $M_h^2 \simeq 2(\lambda_{HS}v_S^2 + \lambda_{HX}v_X^2)$. The mixing angle between the Higgs and the diphoton is experimentally constrained to be small [9, 16]

$$|\sin\theta_{hF}| \lesssim 0.015\sqrt{\Gamma_{\gamma\gamma}/10^{-6}M_F}. \tag{11}$$

In the present model such mixing angle is of order $\theta_{hF} \sim M_h^2 v/M_F^2 v_{S,X}$, which is below its experimental bound provided that $v_{S,X} \gtrsim 500\,\text{GeV}$.

Coming to fermion masses, $\mathcal{U}$ and $\mathcal{U}^c$ split into $N-1$ vector-like up quarks with mass $M_{N-1} = y_X v_X$ and into one with mass $M_1 = \sqrt{(y_S v_S)^2 + (y_X v_X)^2}$, having neglected smaller

electroweak contributions. Recasting LHC searches for similar objects [17], we estimate a bound $M \gtrsim 1.2\ (1.3)\,\mathrm{TeV}$ on their masses for $N = 1\ (5)$.

Diphoton rate

The $\mathcal{U}$ heavy quarks mediate diphoton decays into SM vectors. In the limit where the particles $\wp$ that induce loop decays of a generic scalar φ into SM vectors are much heavier than the energy involved in the decay, their contribution to the decay amplitude is related to their contributions to the β function coefficients $\Delta b_i^\wp$ of the SM gauge couplings as [18]

$$\mathscr{L}_{\text{eff}} = \sum_{i,\wp} \Delta b_i^\wp \frac{\alpha_i}{8\pi} (F_{\mu\nu}^i)^2 \ln \frac{M_\wp(\varphi)}{M_\wp} \tag{12}$$

where $M_\wp(\varphi)$ is the $\wp$ mass for a generic vev of φ. In our model $\varphi = \{x, s\} = \{F, F'\}$ and the loop particles $\wp$ are the quark triplets with mass $M_{N-1} = y_X X$ ($\Delta b_{\text{em}} = 16/9$) and with mass $M_1 = \sqrt{y_X X^2 + y_S |S|^2}$ ($\Delta b_{\text{em}} = (N-1)16/9$). The other Δb_i coefficients are given by eqs. (38 i) and (38k). So

$$\mathscr{L}_{\text{eff}}^\gamma = \frac{\alpha_{\text{em}}}{9\pi} F_{\mu\nu}^2 \left[\sqrt{2} s y_S \frac{\sqrt{M_1^2 - M_{N-1}^2}}{M_1^2} + 2 x y_X \frac{(N-1)M_1^2 + M_{N-1}^2}{M_1^2 M_{N-1}} \right]. \tag{13}$$

Rotating to the mass eigenstates one finds the F width into $\gamma\gamma$:

$$\frac{\Gamma(F \to \gamma\gamma)}{M_F} = \frac{4 M_F^2 \alpha_{\text{em}}^2}{81\pi^3} \left(\frac{\sqrt{2} y_S^2 v_S v_X \cos\theta + 2(N y_X^2 v_X^2 + (N-1) y_S^2 v_S^2) \sin\theta}{v_X (y_S^2 v_S^2 + y_X^2 v_X^2)} \right)^2, \tag{14a}$$

$$\frac{\Gamma(F' \to \gamma\gamma)}{M_{F'}} = \frac{4 M_{F'}^2 \alpha_{\text{em}}^2}{81\pi^3} \left(\frac{\sqrt{2} y_S^2 v_S v_X \sin\theta - 2(N y_X^2 v_X^2 + (N-1) y_S^2 v_S^2) \cos\theta}{v_X (y_S^2 v_S^2 + y_X^2 v_X^2)} \right)^2. \tag{14b}$$

In the limit of small β_{λ_S} the pseudo-Goldstone of scale invariance is the lighter state F:

$$M_F^2 \simeq \frac{2 v_S^2 \beta_{\lambda_S}}{1 + \lambda_{SX}/4\lambda_X} \ll M_{F'}^2 \simeq 4 v_S^2 \lambda_{SX} \left(1 + \frac{\lambda_{SX}}{4\lambda_X}\right), \qquad \tan\theta = \frac{v_X}{\sqrt{2} v_S} \simeq \sqrt{\frac{\lambda_{SX}}{2\lambda_X}} \tag{15}$$

and eq. (14a) reduces to

$$\frac{\Gamma(F \to \gamma\gamma)}{M_F} \simeq \frac{8 N^2 M_F^2 \alpha_{\text{em}}^2}{81\pi^3 (v_S^2 + v_X^2/2)}. \tag{16}$$

In the less relevant opposite limit of small λ_{SX} (and thereby $v_X \ll v_S$) the pseudo-Goldstone of scale invariance is the heavier scalar F':

$$M_F^2 \simeq 4 v_S^2 \lambda_{SX} \ll M_{F'}^2 \simeq 2 v_S^2 \beta_{\lambda_S}, \qquad \tan\theta \simeq \beta_{\lambda_S} \sqrt{\frac{\lambda_X}{\lambda_{SX}^3}} \tag{17}$$

and eq. (14b) reduces to

$$\frac{\Gamma(F' \to \gamma\gamma)}{M_{F'}} \simeq \frac{8 M_{F'}^2 \alpha_{\text{em}}^2}{81\pi^3 v_S^2}. \tag{18}$$



Figure 2: *Running of the couplings for central values for the SM parameters and for* $N = 5$, $g(S_*) \simeq 1.5$, $v_S \simeq 2.2\,\text{TeV}$, $\lambda_{HX}(S_*) \simeq -0.015$, $\lambda_{SX}(S_*) \simeq 0.071$, $\lambda_X(S_*) \simeq 0.030$, $y_S(S_*) = 0.023$ *and* $y_X(S_*) = 0.53$.

Going beyond the approximation of eq. (12) requires computing Feynman diagrams with two different masses in the loop. In the limit $y_S = 0$, such that $M_1 = M_{N-1}$, the full expression is

$$\frac{\Gamma(F \to \gamma\gamma)}{M_F} = \frac{4N^2 M_1^2 \alpha_{\text{em}}^2 y_X^2 \sin^2\theta}{9\pi^3 M_F^2} \left| \mathcal{S}\left(\frac{4M_1^2}{M_F^2}\right) \right|^2 \tag{19}$$

where the loop function $\mathcal{S}$ is

$$\mathcal{S}(x) = 1 + (1-x)\arctan^2\left(\frac{1}{\sqrt{x-1}}\right) \overset{x\gg 1}{\simeq} \frac{2}{3x}. \tag{20}$$

In the limit $M_1 \gg M_F$ eq. (19) reduces to (16).

Numerical example

By performing a global fit of run 1 and run 2 ATLAS and CMS data assuming a narrow width and dominant $gg \to F$ production we find the diphoton rate $\sigma(pp \to F \to \gamma\gamma) = (2.8 \pm 0.7)\,\text{fb}$ at $s = (13\,\text{TeV})^2$, which is reproduced for

$$\frac{\Gamma_{\gamma\gamma}}{M_F} = \frac{s\sigma(pp \to \gamma\gamma)}{K_{gg} C_{gg}} = (3.8 \pm 0.9)\,10^{-7}, \tag{21}$$

where $K_{gg}C_{gg} = 1.5 \times 2140$ are partonic factors [9].

name	spin	$\mathrm{SU}(3)_c$	$\mathrm{SU}(2)_L$	$\mathrm{U}(1)_Y$	$\mathrm{SU}(N)$
X	0	3	1	Y	N
S	0	1	1	0	1
$\mathcal{N}$	1/2	1	1	0	N
$\mathcal{N}^c$	1/2	1	1	0	$\bar{N}$

Table 2: *Beyond the Standard Model field content of the model of type λ) of section 2.2.*

In the limit of small β_{λ_S} the diphoton mass and decay width are reproduced for

$$v_S = N\frac{310\,\mathrm{GeV}}{\sqrt{1+\tan^2\theta}}\sqrt{\frac{10^{-6}M_{\mathcal{F}}}{\Gamma_{\gamma\gamma}}}, \qquad \beta_{\lambda_S} = \frac{5.9}{N^2}(1+\tan^2\theta)^2\frac{\Gamma_{\gamma\gamma}}{10^{-6}M}. \tag{22}$$

Even for $N = 2$ this corresponds to a perturbative value of g, with $Ng^2/(4\pi)^2$ becoming smaller at larger N.

In figure 2 we provide a numerical example with $M_{\mathcal{F}} = 750\,\mathrm{GeV}$, $M_{\mathcal{F}'} \simeq 1.7\,\mathrm{TeV}$, $\Gamma_{\gamma\gamma} \approx 3.5\times10^{-7}M_{\mathcal{F}}$, $M_1 \simeq M_{N-1} \approx 1.3\,\mathrm{TeV}$. There are no Landau poles at energies much smaller than the Planck scale, and the stability conditions of eq. (4) are violated only at low energy, when the desired Coleman-Weinberg mechanism takes place. We assumed central values for the SM parameters, and the Higgs quartic λ_H remains positive up to the Planck scale, unlike in the SM: the new dynamics eliminated the SM vacuum instability [19, 20].[3]

2.2 Model of type λ)

Models of type λ) — those where the diphoton quartic is driven negative by RGE effects of other quartics — need a multiplicity of N scalars in order to avoid Landau poles nearby by sharing the needed relatively large quartics. Some diphoton models introduce a multiplicity of N states for a different reason: in order to mediate a sufficiently large $\mathcal{F} \to \gamma\gamma$ rate. We thereby identify the charged particles that mediate $\mathcal{F} \to \gamma\gamma$ with the scalars that drive the diphoton quartic to negative values.

We now show that a successful model of this type is obtained by considering the massless limit of the diphoton model proposed in [23], where the multiplicity of N states is justified by adding an extra gauge group $\mathrm{SU}(N)$ with gauge coupling g. The model employs the field content listed in table 2. The three scalars are: the SM Higgs doublet H, a neutral singlet S that will contain the diphoton $\mathcal{F}$, and a charged and colored scalar X in the fundamental N of $\mathrm{SU}(N)$, that mediates $\mathcal{F} \to \gamma\gamma, gg$ at one loop.

Including the most generic dimension-less quartic couplings, the scalar potential of the model is:

$$V(H,S,X) \;=\; \lambda_H|H|^4 - \lambda_{HS}|H|^2S^2 + \lambda_S S^4 + \lambda_{HX}|H|^2|X|^2 +$$

[3] Needless to say, new physics at the Planck scale can give new sources of vacuum decay faster than the SM instability scale and faster than the diphoton lifetime. The first part of this statement was strongly emphasised in [21]. For other diphoton models addressing the vacuum instability of the SM see [22].

$$+\lambda_{XS}S^2|X|^2 + \lambda_X \,\mathrm{Tr}(XX^\dagger)^2 + \lambda'_X \,\mathrm{Tr}(XX^\dagger XX^\dagger). \tag{23}$$

The tree-level potential satisfies $V \geq 0$ when the quartic couplings satisfy [15]

$$\begin{cases} \lambda_S > 0, \qquad \lambda_H > 0, \qquad \lambda_X + \alpha\lambda'_X \geq 0, \\ \bar{\lambda}_{HS} \equiv -\lambda_{HS} + 2\sqrt{\lambda_H \lambda_S} \geq 0, \\ \bar{\lambda}_{HX} \equiv \lambda_{HX} + 2\sqrt{\lambda_H(\lambda_X + \alpha\lambda'_X)} \geq 0, \\ \bar{\lambda}_{XS} \equiv \lambda_{XS} + 2\sqrt{\lambda_S(\lambda_X + \alpha\lambda'_X)} \geq 0, \\ \sqrt{\lambda_H}\lambda_{XS} + \sqrt{\lambda_S}\lambda_{HX} - \sqrt{\lambda_X + \alpha\lambda'_X}\lambda_{HS} + 2\sqrt{\lambda_H\lambda_S(\lambda_X + \alpha\lambda'_X)} + \sqrt{\bar{\lambda}_{HS}\bar{\lambda}_{HX}\bar{\lambda}_{XS}} \geq 0, \end{cases} \tag{24}$$

for $\alpha = 1$ and for $\alpha = 1/N_c$ (for $N \geq N_c = 3$), which are the extremal values of

$$\alpha = \,\mathrm{Tr}(XX^\dagger XX^\dagger)/\,\mathrm{Tr}(XX^\dagger)^2.$$

A Coleman-Weinberg minimum is generated when the RGE running of the quartics crosses the boundary of one of these conditions. In practice, we are interested in the case where λ_S becomes negative while running to low energy, while $\lambda_{HS} > 0$ (such that the Higgs too acquires a vev) and $\lambda_{HX} > 0$ (such that X does not acquire any vev).

Furthermore, the model contains N_f extra fermions $\mathcal{N} \oplus \mathcal{N}^c$ with no SM gauge interactions and in the $N \oplus \bar{N}$ representation of $\mathrm{SU}(N)$. Such fermions receive mass from $S\mathcal{N}\mathcal{N}^c$ Yukawa couplings. The lightest among them and among the $\mathrm{SU}(N)$ vectors are Dark Matter candidates. The Yukawa couplings $X^*U\mathcal{N}$ (allowed if $Y = 2/3$) induces X decays into SM up quarks and N. Recasting LHC searches for similar objects [17], we estimate a bound $M_X \gtrsim 1.0\ (1.2)\,\mathrm{TeV}$ on their masses for $N = 3\ (10)$. We assume that such extra Yukawa couplings are small enough that we can neglect their contributions to the RGE. In the presence of these extra fermions, the RGE acquire infra-red fixed points which allow the model to be RGE-extrapolated up to the Planck scale with a stable potential.

The RGE of the model are listed in appendix B. They allow λ_{HS} and λ_{XS} to be naturally small, while $\lambda_{HX} \sim g_1^4$, $\lambda_S \sim g^4$, $\lambda_H \sim g_2^4$.

Masses

The potential at one loop order can be approximated by inserting a running λ_S in the tree-level potential of eq. (24):

$$\lambda_S \simeq \beta_{\lambda_S} \ln \frac{S}{S_*}, \tag{25}$$

where S_* is the scale below which λ_S becomes negative and β_{λ_S} is given in eq. (39a). We can here neglect the running of the other couplings. Expanding the scalars as

$$S = w + s, \qquad H = \frac{1}{\sqrt{2}}\begin{pmatrix} 0 \\ v + h \end{pmatrix} \tag{26}$$

the effective potential is minimised by

$$w = S_* \exp\left[-\frac{1}{4} + \frac{\lambda_{HS}^2}{4\lambda_H \beta_{\lambda_S}}\right], \qquad v = w\sqrt{\frac{\lambda_{HS}}{\lambda_H}}. \tag{27}$$

The scalar mass matrix at the minimum in the (h, s) basis is

$$2v^2 \begin{pmatrix} \lambda_H & -\sqrt{\lambda_H \lambda_{HS}} \\ -\sqrt{\lambda_H \lambda_{HS}} & \lambda_{HS} + 2\beta_{\lambda_S} \lambda_H / \lambda_{HS} \end{pmatrix}. \tag{28}$$

The mass eigenstates are the physical Higgs and the diphoton F. In the limit of small $\epsilon \equiv \lambda_{HS}^2 / 2\lambda_H \beta_{\lambda_S}$, which corresponds to a Higgs mass smaller than the diphoton mass, the mass eigenvalues are

$$M_h^2 \simeq 2v^2 \lambda_H (1 - \epsilon + \cdots), \qquad M_F^2 \simeq 4w^2 \beta_{\lambda_S} (1 + \epsilon + \cdots). \tag{29}$$

The diphoton mass M_F is suppressed by a one loop factor because the diphoton is identified here with the pseudo-Goldstone boson of scale invariance. The h/F mixing angle

$$\theta_{hF} \simeq \frac{M_h^2}{M_F^2} \frac{v}{w} \tag{30}$$

is below the experimental bound of eq. (11) for $w \gtrsim 500\,\mathrm{GeV}$. Finally, X acquires the mass $M_X \simeq w\sqrt{\lambda_{XS}}$, the extra fermions receive mass $M_{\mathcal{N}} = y_S w$ from $y_S\, S\mathcal{N}\mathcal{N}^c$ Yukawa couplings, and the SM particles acquire the usual masses through the Higgs vacuum expectation value.

Diphoton rate

The $\Gamma_{\gamma\gamma} \equiv \Gamma(F \to \gamma\gamma)$ rate is given by

$$\frac{\Gamma_{\gamma\gamma}}{M_F} = \frac{9N^2 \alpha_{\mathrm{em}}^2 Y^4}{256\pi^3} \left| \frac{w \lambda_{XS} M_F}{M_X^2} F\left(\frac{4M_X^2}{M_F^2}\right) \right|^2 \tag{31}$$

where the loop function F is

$$F(x) = x\left[x \arctan^2\left(\frac{1}{\sqrt{x-1}}\right) - 1 \right] \overset{x\to\infty}{=} \frac{1}{3}. \tag{32}$$

In the limit $M_X \gg M_F/2$ the $\Gamma_{\gamma\gamma}$ rate does not depend on λ_{HS}, as can be understood using the Low Energy Theorem of eq. (12), taking into account that, in the present model, $\wp = X$ with $M_X(F)/M_X = F/w + \cdots$.

As usual a large multiplicity N enhances $\Gamma_{\gamma\gamma}$; in our context it also enhances M_F. In order to reproduce desired values of $M_h = 125\,\mathrm{GeV}$, $M_F = 750\,\mathrm{GeV}$, M_X and $\Gamma_{\gamma\gamma}$, the model parameters are fixed to the following values

$$\begin{aligned} \lambda_{XS} &\approx 0.24 Y^4 \left(\frac{\mathrm{TeV}}{M_X}\right)^6 \left(\frac{10^{-6}}{\Gamma_{\gamma\gamma}/M_F}\right), & w &\approx \frac{2.0\,\mathrm{TeV}}{Y^2} \left(\frac{M_X}{\mathrm{TeV}}\right)^4 \left(\frac{\Gamma_{\gamma\gamma}/M_F}{10^{-6}}\right)^{1/2}, \\ \lambda_{HS} &\approx 0.002\, Y^4 \left(\frac{10^{-6}}{\Gamma_{\gamma\gamma}/M_F}\right), & N &\approx \frac{30}{Y^4} \left(\frac{M_X}{\mathrm{TeV}}\right)^4 \left(\frac{\Gamma_{\gamma\gamma}/M_F}{10^{-6}}\right). \end{aligned} \tag{33}$$

at leading order in $M_h/M_F \ll 1$, and neglecting higher order corrections such as the running of the coupling constants between M_F and S_*. We see that N is large (tens) for $Y = 2/3$ and depends strongly on Y, such that a small N is obtained for $Y = 4/3$ or $5/3$.



Figure 3: *Running of the couplings for central values for the SM parameters, that we took from [19], and* $g^2(M_F) = (4\pi)^2/N$. *We find that that varying the values of* $\lambda_X(M_F)$, $\lambda'_X(M_F)$, $\lambda_{HX}(M_F)$ *have a very small impact on the running at energies much bigger than* F_*. **Upper plots:** *We assumed* $Y = 2/3$, $N = 12$, $w = 1890\,\mathrm{GeV}$, $5N$ *fermions in the* $N \oplus \bar{N}$ *representation of* $\mathrm{SU}(N)$*; we also take* $\lambda_X(M_F) = 0.15$, $\lambda'_X(M_F) = 0.28$, $\lambda_{HX}(M_F) = 0.17$. **Lower plots:** *We assumed* $Y = 4/3$, $N = 5$, $w = 2000\,\mathrm{GeV}$, *26 fermions in the* $N \oplus \bar{N}$ *representation of* $\mathrm{SU}(N)$*; we also take* $\lambda_X(M_F) = 0.15$, $\lambda'_X(M_F) = 0.28$, $\lambda_{HX}(M_F) = 0.47$.

name	spin	$\mathrm{SU}(3)_c$	$\mathrm{SU}(2)_L$	$\mathrm{U}(1)_Y$	$\mathrm{SU}(N)$
$\mathcal{Q}$	$1/2$	3	2	$+1/6$	N
$\mathcal{Q}^c$	$1/2$	$\bar{3}$	2	$-1/6$	$\bar{N}$
$\mathcal{U}$	$1/2$	$\bar{3}$	1	$-2/3$	N
$\mathcal{U}^c$	$1/2$	3	1	$+2/3$	$\bar{N}$

Table 3: *Beyond the Standard Model field content of the strongly-coupled model of section 3.1.*

Numerical examples

In figure 3 we provide two numerical examples with $M_{\mathcal{F}} = 750\,\mathrm{GeV}$, with no Landau poles at energies much smaller than the Planck scale and with all the stability conditions of eq. (24) satisfied. We assumed central values for the SM parameters, and the Higgs quartic λ_H remains positive up to the Planck scale unlike in the SM.

In the upper example we have $Y = 2/3$ and $M_X \approx 1.2\,\mathrm{TeV}$ The example leads to $\Gamma_{\gamma\gamma}/M_{\mathcal{F}} \simeq 3.1 \times 10^{-7}$ and to a diphoton decay rate in two gluons $\Gamma_{gg}/M_{\mathcal{F}} \simeq 4.7 \times 10^{-5}$. This example employs $N = 12$, which is compatible with experimental bounds.

Given that such a large N might look implausible from a low-energy perspective, we provide in the lower row a second example with $N = 5$, achieved by increasing $Y = 4/3$. This example has $M_X = 1.6\,\mathrm{TeV}$, $\Gamma_{\gamma\gamma} \simeq 4.4 \times 10^{-7} M_{\mathcal{F}}$, $\Gamma_{gg} \simeq 4.2 \times 10^{-6} M_{\mathcal{F}}$.

In both cases the $\mathrm{SU}(N)$ gauge constant becomes relatively large at the diphoton scale such that the couplings (in particular λ_{XS}) have a fast running; we tried to include such corrections by renormalising couplings at appropriate scales.

3 Strongly coupled models

In this section we try to build dimensionless models where a new gauge interaction (we use for it the old-fashioned name TechniColor, or TC) becomes strong around the weak scale, inducing the weak scale and the diphoton $\mathcal{F}$. Like in the weakly-coupled case, it is easy to reproduce the diphoton excess by adding one extra ad-hoc diphoton scalar (see [24] for one such model). We are interested in models where the diphoton automatically emerges as a bound state of the TC dynamics. In view of strong LHC bounds on extra bound states, especially the ones with color, plausible models identify the diphoton with a bound state that is much lighter than the others. The diphoton could be a TCη (here discussed in section 3.1) or a TC-dilaton (here discussed in section 3.2). The TC-dilaton is especially interesting from our point of view, given that it is the pseudo-Goldstone boson of scale invariance.

3.1 The diphoton as a TCη

A class of strongly-coupled models that aims at reproducing the diphoton excess are those where the diphoton $\mathcal{F}$ is identified with a TCη composite pseudo-scalar, given that this field is a light pseudo-Goldstone boson with anomalous couplings to SM vectors. The various

models proposed in the literature [9, 25] employ massive techni-quarks, and a massive SM Higgs doublet. The masses are assumed to be comparable to the TC scale, but the coincidence is left unexplained.

Our goal is exploring whether such masses can be all set to zero, obtaining a dimensionless model. For concreteness, we consider a model with $G_{\rm TC} = {\rm SU}(N)$ and the techni-quark content of table 3[4] which allows for two Yukawa couplings to the SM Higgs doublet H:

$$y_1 H\mathcal{Q}\mathcal{U} + y_2 H^\dagger \mathcal{Q}^c\mathcal{U}^c + \text{h.c.} = H\bar{\Psi}_{\mathcal{Q}}(y + i\gamma_5\tilde{y})\Psi_{\mathcal{U}} + \text{h.c.} \tag{34}$$

In the latter expression we introduced Dirac spinors Ψ and the scalar coupling $y = (y_1 + y_2^*)/2$ and the pseudo-scalar coupling $\tilde{y} = i(y_1 - y_2^*)/2$, such that $|y| \ll |\tilde{y}|$ and $|\tilde{y}| \ll |y|$ are radiatively stable special cases, that can be justified by assuming a CP-like symmetry.

We now discuss the composite states. Among the many states around the TCρ mass, $m_\rho \sim g_\rho f_{\rm TC}$ with $g_\rho \sim 4\pi/\sqrt{N}$, there is the TC $\eta' \sim QQ^c + UU^c$ singlet which receives a mass from TC anomalies. The TC dynamics breaks the global accidental symmetry ${\rm SU}(9)_L \otimes {\rm SU}(9)_R \to {\rm SU}(9)_V$ giving 80 lighter techni-pions, with the following SM quantum numbers:

$$(1,1)_0 \oplus (1,2)_{\pm 1/2} \oplus (1,3)_0 \oplus 2(8,1)_0 \oplus (8,2)_{\pm 1/2} \oplus (8,3)_0. \tag{35}$$

Taking into account that we assume massless techni-quarks, the techni-pions consist of:

72) 9 color octets, QQ^c, UU^c, QU^c, Q^cU, which get positive squared masses at loop level from QCD interactions, $m \approx \sqrt{\frac{3}{4\pi}\alpha_3 C m_\rho^2} \sim 0.2 m_\rho$, where $C = (N^2-1)/2N$;

3) a ${\rm SU}(2)_L$ triplet QQ^c, which similarly gets a positive squared mass from weak interactions;

1) a TC $\eta \sim UU^c - \frac{1}{2}QQ^c$ singlet (to be identified with the diphoton), which gets a squared mass from Yukawa loop corrections with unknown sign and of order $m_\eta \sim y m_\rho/4\pi$, see appendix A.3 of [6];

4) a TCpion π_2 from QU^c, Q^cU with the same gauge quantum numbers as the Higgs doublet forms whenever gauge quantum numbers allow for a Yukawa coupling to H. It gets a mass $m_{\pi_2} \sim g_2 m_\rho/4\pi$ from weak gauge interactions, plus another contribution from $y_{1,2}$.

Furthermore, π_2 acquires a tree level mixing with the Higgs boson. The resulting mass matrix is

$$\begin{array}{c} \\ \pi_2 \\ H \end{array} \begin{array}{c} \begin{array}{cc} \pi_2^* & \quad H^* \end{array} \\ \begin{pmatrix} (\mathcal{O}(g_2^2) \pm \mathcal{O}(y^2))/(4\pi)^2 & \mathcal{O}(y)\sqrt{N}/(4\pi) \\ \mathcal{O}(y)\sqrt{N}/(4\pi) & -\mathcal{O}(y^2)N/(4\pi)^2 \end{pmatrix} \end{array} m_\rho^2 \tag{36}$$

The tree-level mixing induces a negative see-saw contribution $\Delta m_H^2 \sim -\tilde{y}^2 m_\rho^2 f_{\rm TC}^2/m_{\pi_2}^2$ to the Higgs mass parameter, where . This contribution is too big, given that the diphoton only gets a mass at loop level.

[4] A similar model with up-type quarks replaced by down-type quarks has been considered in [26] because the lightest TCbaryon is a good Dark Matter candidate.



Figure 4: *Correction to the Higgs and diphoton mass coming from the Yukawa couplings.*

Such tree-level contribution vanishes if we assume $\tilde{y} = 0$ – a natural special case that respects a CP-like parity. Indeed, H is a scalar, while π_2 and η are pseudo-scalars. Their tree-level potential includes a CP-conserving term $\sim y m_\rho H \pi_2^* \eta + \text{h.c.}$ As a result, both H and of η receive loop-level masses of order $y m_\rho / 4\pi$, with no symmetry relation among them, given that H is elementary while η is composite, see fig. 4. With a relatively large value of $y \sim$ few the model can give $M_{\not F} \approx 750\,\text{GeV}$ together with $f_{\rm TC} \approx 100\,\text{GeV} \times N$ as demanded by the diphoton rate suggested by preliminary ATLAS and CMS results. The η eigenstate has a vanishing color anomaly (see table 2 of [27]); this is not a problem because, taking into loop corrections, the mass eigenstate $\not F \approx \eta + \eta' y^2/(4\pi)^2$ acquires a sufficiently large coupling to gluons.

3.2 The diphoton as a TC-dilaton or TCσ

In strongly coupled TC models with a QCD-like dynamics the mass of the pseudo-Goldstone boson of scale invariance is not suppressed with respect to the other composite states, because strong interactions give a fast running that strongly breaks scale invariance. On the other hand, such state is somehow lighter than the other bound states in models with a 'walking' dynamics (namely, a β function of the new gauge interaction which remains somewhat small), which is obtained in TC models with a larger matter content than QCD. This lightness is beneficial for the diphoton phenomenology, given that other colored bound states must be heavier than $750\,\text{GeV}$ in order to satisfy LHC bounds.

From a low-energy perspective, such light state is the σ field sometimes explicitly included in effective chiral Lagrangians, where its vacuum expectation value, $\sigma = f_{\rm TC} + \not F$, breaks scale invariance as well as a global chiral symmetry, such that $f_{\rm TC}$ becomes the techni-pion decay constant. See [28] for a recent discussion of the σ, and [9, 29] for recent discussions of the dilaton. In the limit where $\not F$ is lighter than the other bound states, its coupling to SM vectors is dictated by eq. (12): using $\ln M_\wp(\not F)/M_\wp \simeq \not F / f_{\rm TC}$ we find

$$\frac{\Gamma_{\gamma\gamma}}{M_{\not F}} = 10^{-6} \left(\frac{120\,\text{GeV}}{f_{\rm TC}} \right)^2 \Delta b_{\rm em}^2 \tag{37}$$

where $\Delta b_{\rm em}$ is the techniquark contribution to the running of the electromagnetic coupling. For example, a color triplet techni-fermion with hypercharge Y (such as $\mathcal{U} \oplus \mathcal{U}^c$ in table 1) contributes as $\Delta b_{\rm em} = 4NY^2$, while a techni-scalar contributes 4 times less. The SM $\Delta b_{Y,2,3}$ must be smaller than about 10, in order to avoid sub-Planckian Landau poles for the SM gauge couplings.

The dynamically generated Higgs mass depends on the model. In the minimal case where the Higgs has no direct coupling to techni-particles, electro-weak loop effects induce a contribution to its squared mass of order $M_h^2 \sim -\alpha_Y^2 f_{\rm TC}^2$ [6], which is negative but small, in view of eq. (37). A larger model-dependent tree-level contribution is obtained if the Higgs has a Yukawa coupling to techni-fermions or a quartic coupling to techni-scalars.

4 Conclusions

We proposed dimension-less models where a new resonance F decaying into two photons dynamically breaks scale invariance, generating a weak scale hierarchically smaller than the Planck scale. The diphoton channel is interesting as it is particularly clean and has therefore a great discovery potential. All particles acquire their masses from their couplings to F: thereby the smoking gun of this scenario is observing that F couples to all particles proportionally to their mass.

As a benchmark case we identified F with the 750 GeV resonance hinted by LHC data. Although it could very well be that this excess is a statistical fluctuation, it nevertheless provides an interesting example. The $F \to \gamma\gamma$ rate suggested by LHC data is obtained adding extra charged particles, heavier than the SM particles and thereby more strongly coupled to F.

Diphoton models of this type generically need that such extra particles have masses around the weak scale. Unlike the SM fermions, such extra particles have no chirality reason to be around the weak scale. Scale invariance provides one possible reason: like the Higgs, these extra charged particles acquire a mass from F. All particles are massless until the diphoton develops a vev or condensate w.

A generic scalar that acts as the 'Higgs of the Higgs' can have a w and/or a mass M much larger than the weak scale, provided that it is very weakly coupled to the Standard Model (SM). This is not possible if such scalar is identified with the diphoton: w and M are now fixed by the diphoton mass and rate in $\gamma\gamma$.

In section 2.1 we presented a weakly-coupled dimension-less model where the diphoton is charged under extra gauge interactions, which induce the Coleman-Weinberg mechanism. In section 2.2 we presented a weakly-coupled dimension-less model where the Coleman-Weinberg mechanism is induced by quartic interactions of the diphoton with the charged scalars that mediate $F \to \gamma\gamma$.

Both models can be RGE-extrapolated up to the Planck scale, thereby generating the large hierarchy with respect to the weak scale. Both models contain Dark Matter candidates. Both models remove the instability of the SM potential. Both models give rise to an extended

phase transition when the diphoton and the Higgs acquire vacuum expectation values. Such phase transition can be of first order, possibly giving gravitational wave signals [30] and the baryon asymmetry [31].

In section 3.1 we discussed dimension-less strongly-coupled models where the diphoton is a pseudo-scalar bound state analogous of the η in QCD. In section 3.2 we discussed dimension-less strongly-coupled models where the diphoton is a scalar bound state analogous of the σ in chiral effective Lagrangians, or of a dirty dilaton in fundamental strongly coupled models with walking dynamics. Both scenarios can produce a weak scale lighter than the diphoton mass by $M_h/M_{\Gamma} \sim 1/6$, but this needs extra model building features.

Acknowledgements

This work was supported by the grant 669668 – NEO-NAT – ERC-AdG-2014. Kristjan Kannike was supported by the Estonian Research Council grant PUT799, the grant IUT23-6 of the Estonian Ministry of Education and Research, and by the EU through the ERDF CoE program. We thank Roberto Franceschini and Francesco Riva for useful discussions.

A RGE for model *g*)

Defining $\beta_g = dg/d\ln\mu$, the one-loop RGE for the new couplings are

$$
\begin{aligned}
(4\pi)^2\beta_{\lambda_S} &= 4(4+N)\lambda_S^2 + \frac{3(N-1)(N^2+2N-2)}{4N^2}g^4 + \\
&\quad +2(\lambda_{SX}^2+\lambda_{HS}^2) - 6y_S^4 + \lambda_S\left[12y_S^2 - 6\frac{N^2-1}{N}g^2\right], \\
(4\pi^2)\beta_{\lambda_{HS}} &= 12y_S^2y_t^2 - 4\lambda_{HS}^2 - 4\lambda_{HX}\lambda_{SX} + \qquad (38a)\\
&\quad +\lambda_{HS}\left[-3g^2\frac{N^2-1}{N} - \frac{9g_1^2}{10} - \frac{9g_2^2}{2} + 6y_S^2 + 6y_t^2 + 12\lambda_H + (4N+4)\lambda_S\right], \\
(4\pi^2)\beta_{\lambda_{HX}} &= -8\lambda_{HX}^2 - 2N\lambda_{HS}\lambda_{SX} + \qquad (38b)\\
&\quad +\lambda_{HX}\left[-\frac{9g_1^2}{10} - \frac{9g_2^2}{2} + 6y_t^2 + 12Ny_X^2 + 12\lambda_H + 24\lambda_X\right], \qquad (38c)\\
(4\pi^2)\beta_{\lambda_{SX}} &= 12y_S^2y_X^2 - 8\lambda_{SX}^2 - 4\lambda_{HS}\lambda_{HX} + \\
&\quad +\lambda_{SX}\left[-3g^2\frac{N^2-1}{N} + 6y_S^2 + 12Ny_X^2 + (4N+4)\lambda_S + 24\lambda_X\right], \qquad (38d)\\
(4\pi^2)\beta_{\lambda_X} &= -6Ny_X^4 + 24N\lambda_X y_X^2 + 2\lambda_{HX}^2 + N\lambda_{SX}^2 + 72\lambda_X^2, \qquad (38e)\\
(4\pi^2)\beta_{y_X} &= (6N+3)y_X^3 + y_X\left[-3\frac{N^2-1}{N}g^2 - \frac{8g_1^2}{5} - 8g_3^2 + \frac{y_S^2}{2}\right], \qquad (38f)\\
(4\pi^2)\beta_{y_S} &= \frac{N+7}{2}y_S^3 + y_S\left[-\frac{3}{2}g^2\frac{N^2-1}{N} - \frac{8g_1^2}{5} - 8g_3^2 + y_t^2 + \frac{y_X^2}{2}\right], \qquad (38g)\\
(4\pi^2)\beta_g &= \left(\frac{13}{6} - \frac{11N}{3}\right)g^3. \qquad (38h)
\end{aligned}
$$

Finally, the RGE for the SM couplings are

$$(4\pi^2)\beta_{g_1} = \left(\frac{41}{10} + \Delta b_1\right) g_1^3, \qquad \Delta b_1 = \frac{16N}{15}, \tag{38i}$$

$$(4\pi^2)\beta_{g_2} = -\frac{19 g_2^3}{6}, \tag{38j}$$

$$(4\pi^2)\beta_{g_3} = (-7 + \Delta b_3)\, g_3^3, \qquad \Delta b_3 = \frac{2N}{3}, \tag{38k}$$

$$(4\pi^2)\beta_{y_t} = \frac{9y_t^3}{2} + y_t\left[-\frac{17g_1^2}{20} - \frac{9g_2^2}{4} - 8g_3^2 + \frac{N y_S^2}{2}\right], \tag{38l}$$

$$(4\pi^2)\beta_{\lambda_H} = \frac{27g_1^4}{200} + \frac{9g_2^2 g_1^2}{20} + \frac{9g_2^4}{8} - 6y_t^4 + 24\lambda_H^2 + N\lambda_{HS}^2 + 2\lambda_{HX}^2 + \tag{38m}$$

$$+\lambda_H\left[12y_t^2 - \frac{9g_1^2}{5} - 9g_2^2\right]. \tag{38n}$$

B RGE for model λ)

Defining $\beta_g = dg/d\ln\mu$, the RGE for the new couplings are

$$(4\pi)^2\beta_{\lambda_S} = 72\lambda_S^2 + 3N\lambda_{XS}^2 + 2\lambda_{HS}^2, \tag{39a}$$

$$(4\pi)^2\beta_{\lambda_X} = 4(3N+4)\lambda_X^2 + 12\lambda_X'^2 + 2\lambda_{XS}^2 + 2\lambda_{HX}^2 + \tag{39b}$$

$$+\lambda_X\left[8(3+N)\lambda_X' - \frac{6(N^2-1)}{N}g^2 - 16g_3^2 - \frac{36Y^2g_1^2}{5} + \frac{54Y^4}{25}g_1^4\right] +$$

$$+\frac{3(N^2+2)}{4N^2}g^4 + \frac{11}{12}g_3^4 + \frac{3N+1}{N}g^2g_3^2 - \frac{18Y^2g^2g_1^2}{5N} - \frac{6Y^2g_3^2g_1^2}{5},$$

$$(4\pi)^2\beta_{\lambda_X'} = 4(3+N)\lambda_X'^2 + \lambda_X'\left[24\lambda_X - \frac{36Y^2g_1^2}{5} - \frac{6(N^2-1)}{N}g^2 - 16g_3^2\right] + \tag{39c}$$

$$+\frac{3(N^2-4)}{4N}g^4 + \frac{5g_3^4}{4} - \frac{N+3}{N}g^2g_3^2 + \frac{18Y^2g^2g_1^2}{5} + \frac{18Y^2g_3^2g_1^2}{5},$$

$$(4\pi)^2\beta_{\lambda_{XS}} = 8\lambda_{XS}^2 - 4\lambda_{HS}\lambda_{HX} + 4\lambda_{XS}\Big[(1+3N)\lambda_X + (3+N)\lambda_X' + \tag{39d}$$

$$+6\lambda_S - \frac{9Y^2g_1^2}{10} - \frac{3(N^2-1)}{4N}g^2 - 2g_3^2\Big],$$

$$(4\pi)^2\beta_{\lambda_{HX}} = \lambda_{HX}\left[4(1+3N)\lambda_X + 4(3+N)\lambda_X' - \frac{(36Y^2+9)g_1^2}{10} - \frac{9g_2^2}{2} + 12\lambda_H + 6y_t^2\right] + \tag{39e}$$

$$-4\lambda_{HS}\lambda_{XS} + 4\lambda_{HX}^2 + \frac{27g_1^4Y^2}{25} - \frac{3(N^2-1)}{N}\lambda_{HX}g^2 - 8\lambda_{HX}g_3^2,$$

$$(4\pi)^2\beta_{\lambda_{HS}} = \lambda_{HS}\left[24\lambda_S - \frac{9g_1^2}{10} - \frac{9g_2^2}{2} + 6y_t^2 + 12\lambda_H\right] - 6N\lambda_{XS}\lambda_{HX} - 8\lambda_{HS}^2, \tag{39f}$$

$$(4\pi)^2\beta_g = g^3\left(-\frac{11}{3}N + \frac{1}{2} + \frac{2}{3}N_f\right). \tag{39g}$$

Finally, the RGE for the SM couplings are

$$(4\pi)^2\beta_{g_1} = g_1^3\frac{41 + 6NY^2}{10}, \tag{39h}$$

$$(4\pi)^2\beta_{g_2} = -\frac{19g_2^3}{6}, \tag{39i}$$

$$(4\pi)^2\beta_{g_3} = g_3^3\left(-7+\frac{N}{6}\right), \tag{39j}$$

$$\begin{aligned}(4\pi)^2\beta_{\lambda_H} = {} & 2\lambda_{HS}^2+3N\lambda_{HX}^2+\frac{27g_1^4}{200}+\frac{9g_1^2g_2^2}{20}+\frac{9g_2^4}{8}+\\ & +\lambda_H\left(-\frac{9g_1^2}{5}-9g_2^2+12y_t^2\right)+24\lambda_H^2-6y_t^4,\end{aligned} \tag{39k}$$

$$(4\pi)^2\beta_{y_t} = y_t\left(\frac{9}{2}y_t^2-\frac{17g_1^2}{20}-8g_3^2-\frac{9g_2^2}{4}\right). \tag{39l}$$

References

[1] M. Farina, D. Pappadopulo, A. Strumia, *"A modified naturalness principle and its experimental tests"*, JHEP 1308 (2013) 022 [arXiv:1303.7244]. M. Heikinheimo, A. Racioppi, M. Raidal, C. Spethmann, K. Tuominen, *"Physical Naturalness and Dynamical Breaking of Classical Scale Invariance"*, Mod. Phys. Lett. A29 (2014) 1450077 [arXiv:1304.7006].

[2] S. L. Adler, *"A Formula for the Induced Gravitational Constant"*, Phys. Lett. B **95** (1980) 241. A. Zee, *"Spontaneously Generated Gravity"*, Phys. Rev. D 23 (1981) 858. A. Salvio, A. Strumia, *"Agravity"*, JHEP 1406 (2014) 080 [arXiv:1403.4226]. M. Rinaldi, G. Cognola, L. Vanzo, S. Zerbini, *"Inflation in scale-invariant theories of gravity"*, Phys. Rev. D91 (2015) 123527 [arXiv:1410.0631]. K. Kannike, G. Htsi, L. Pizza, A. Racioppi, M. Raidal, A. Salvio, A. Strumia, *"Dynamically Induced Planck Scale and Inflation"*, JHEP 1505 (2015) 065 [arXiv:1502.01334]. A. Farzinnia, S. Kouwn, *"Classically scale invariant inflation, supermassive WIMPs, and adimensional gravity"*, Phys. Rev. D93 (2016) 063528 [arXiv:1512.05890]. A. Farzinnia, *"Classically Scale Invariant Inflation and (A)gravity"* [arXiv:1512.08727]. A. Salvio, A. Strumia, *"Quantum mechanics of 4-derivative theories"*, Eur. Phys. J. C76 (2016) 227 [arXiv:1512.01237]. M.B. Einhorn, D.R.T. Jones, *"Induced Gravity II: Grand Unification"* [arXiv:1602.06290].

[3] T. Hambye, A. Strumia, *"Dynamical generation of the weak and Dark Matter scale"*, Phys. Rev. D88 (2013) 055022 [arXiv:1306.2329].

[4] R. Foot, A. Kobakhidze, K.L. McDonald, R.R. Volkas, *"Neutrino mass in radiatively-broken scale-invariant models"*, Phys. Rev. D76 (2007) 075014 [arXiv:0706.1829] R. Foot, A. Kobakhidze, K.L. McDonald, R.R. Volkas, *"A Solution to the hierarchy problem from an almost decoupled hidden sector within a classically scale invariant theory"*, Phys. Rev. D77 (2007) 035006 [arXiv:0709.2750]. R. Foot, A. Kobakhidze, R.R. Volkas, *"Stable mass hierarchies and dark matter from hidden sectors in the scale-invariant standard model"*, Phys. Rev. D82 (2010) 035005 [arXiv:1006.0131]. R. Foot, A. Kobakhidze, *"Electroweak Scale Invariant Models with Small Cosmological Constant"*, Int. J. Mod. Phys. A30 (2015) 1550126 [arXiv:1112.0607]. L. Alexander-Nunneley, A. Pilaftsis, *"The Minimal Scale Invariant Extension of the Standard Model"*, JHEP 1009 (2010) 021 [arXiv:1006.5916]. K. Ishiwata, *"Dark Matter in Classically Scale-Invariant Two Singlets Standard Model"*, Phys. Lett. B710 (2012) 134 [arXiv:1112.2696]. J.S. Lee, A. Pilaftsis, *"Radiative Corrections to Scalar Masses and Mixing in a Scale Invariant Two Higgs Doublet Model"*, Phys. Rev. D86 (2012) 035004 [arXiv:1201.4891]. A. Farzinnia, H-J. He, J. Ren,

"Natural Electroweak Symmetry Breaking from Scale Invariant Higgs Mechanism", Phys. Lett. B727 (2013) 141 [arXiv:1308.0295]. D. Chway, T.H. Jung, H.D. Kim, R. Dermisek, *"Radiative Electroweak Symmetry Breaking Model Perturbative All the Way to the Planck Scale"*, Phys. Rev. Lett. 113 (2014) 051801 [arXiv:1308.0891]. E. Gabrielli, M. Heikinheimo, K. Kannike, A. Racioppi, M. Raidal, C. Spethmann, *"Towards Completing the Standard Model: Vacuum Stability, EWSB and Dark Matter"*, Phys. Rev. D89 (2014) 015017 [arXiv:1309.6632]. C.T. Hill, *"Is the Higgs Boson Associated with Coleman-Weinberg Dynamical Symmetry Breaking?"*, Phys. Rev. D89 (2014) 073003 [arXiv:1401.4185] J. Guo, Z. Kang, *"Higgs Naturalness and Dark Matter Stability by Scale Invariance"*, Nucl. Phys. B898 (2015) 415 [arXiv:1401.5609]. H. Davoudiasl, I.M. Lewis, *"Right-Handed Neutrinos as the Origin of the Electroweak Scale"*, Phys. Rev. D90 (2014) 033003 [arXiv:1404.6260] K. Allison, C.T. Hill, G.G. Ross, *"Ultra-weak sector, Higgs boson mass, and the dilaton"*, Phys. Lett. B738 (2014) 191 [arXiv:1404.6268]. A. Farzinnia, J. Ren, *"Higgs Partner Searches and Dark Matter Phenomenology in a Classically Scale Invariant Higgs Boson Sector"*, Phys. Rev. D90 (2014) 015019 [arXiv:1405.0498]. M. Lindner, S. Schmidt, J. Smirnov, *"Neutrino Masses and Conformal Electro-Weak Symmetry Breaking"*, JHEP 1410 (2014) 177 [arXiv:1405.6204]. K. Allison, C.T. Hill, G.G. Ross, *"An ultra-weak sector, the strong CP problem and the pseudo-Goldstone dilaton"*, Nucl. Phys. B891 (2014) 613 [arXiv:1409.4029] K. Endo, Y. Sumino, *"A Scale-invariant Higgs Sector and Structure of the Vacuum"*, JHEP 1505 (2015) 030 [arXiv:1503.02819]. A. Karam, K. Tamvakis, *"Dark matter and neutrino masses from a scale-invariant multi-Higgs portal"*, Phys. Rev. D92 (2015) 075010 [arXiv:1508.03031]. A. Ahriche, K.L. McDonald, S. Nasri, *"The Scale-Invariant Scotogenic Model"* [arXiv:1604.05569]

[5] S. Benic, B. Radovcic, *"Electroweak breaking and Dark Matter from the common scale"*, Phys. Lett. B732 (2014) 91 [arXiv:1401.8183]. G.M. Pelaggi, *"Predictions of a model of weak scale from dynamical breaking of scale invariance"*, Nucl. Phys. B893 (2015) 443 [arXiv:1406.4104]. W. Altmannshofer, W.A. Bardeen, M. Bauer, M. Carena, J.D. Lykken, *"Light Dark Matter, Naturalness, and the Radiative Origin of the Electroweak Scale"*, JHEP 1501 (2015) 032 [arXiv:1408.3429]. A.D. Plascencia, *"Classical scale invariance in the inert doublet model"*, JHEP 1509 (2015) 026 [arXiv:1507.04996]. N. Haba, H. Ishida, N. Okada, Y. Yamaguchi, *"Bosonic seesaw mechanism in a classically conformal extension of the Standard Model"*, Phys. Lett. B754 (2016) 349 [arXiv:1508.06828]. A. Karam, K. Tamvakis, *"Dark matter and neutrino masses from a classically scale-invariant multi-Higgs portal"* [arXiv:1603.08470].

[6] O. Antipin, M. Redi, A. Strumia, *"Dynamical generation of the weak and Dark Matter scales from strong interactions"*, JHEP 1501 (2015) 157 [arXiv:1410.1817].

[7] M. Holthausen, J. Kubo, K.S. Lim, M. Lindner, *"Electroweak and Conformal Symmetry Breaking by a Strongly Coupled Hidden Sector"*, JHEP 1312 (2013) 076 [arXiv:1310.4423]. J. Kubo, K.S. Lim, M. Lindner, *"Electroweak Symmetry Breaking via QCD"*, Phys. Rev. Lett. 113 (2014) 091604 [arXiv:1403.4262]. C.D. Carone, R. Ramos, *"Dark chiral symmetry breaking and the origin of the electroweak scale"*, Phys. Lett. B746 (2015) 424 [arXiv:1505.04448]. H. Ishida, S. Matsuzaki, Y. Yamaguchi, *"Invisible Axion-Like Dark Matter from Electroweak Bosonic Seesaw"* [arXiv:1604.07712].

[8] ATLAS note, ATLAS-CONF-2015-081, *"Search for resonances decaying to photon pairs in 3.2 fb^{-1} of pp collisions at $\sqrt{s} = 13\,\mathrm{TeV}$ with the ATLAS detector"*. CMS note, CMS PAS EXO-15-004 *"Search for new physics in high mass diphoton events in proton-proton collisions at $13\,\mathrm{TeV}$"*.

[9] R. Franceschini, G.F. Giudice, J.F. Kamenik, M. McCullough, A. Pomarol, R. Rattazzi, M. Redi, F. Riva, A. Strumia, R. Torre, *"What is the $\gamma\gamma$ resonance at $750\,\mathrm{GeV}$?"*, JHEP 1603 (2016) 144 [arXiv:1512.04933].

[10] S. Di Chiara, L. Marzola, M. Raidal, *"First interpretation of the 750GeV diphoton resonance at the LHC"*, Phys. Rev. D93 (2016) 095018 [arXiv:1512.04939]. C. Petersson, R. Torre, *"$750\,\mathrm{GeV}$ Diphoton Excess from the Goldstino Superpartner"*, Phys. Rev. Lett. 116 (2016) 151804 [arXiv:1512.05333]. S.V. Demidov, D.S. Gorbunov, *"On sgoldstino interpretation of the diphoton excess"* [arXiv:1512.05723]. T-F. Feng, X-Q. Li, H-B. Zhang, S-M. Zhao, *"The LHC 750 GeV diphoton excess in supersymmetry with gauged baryon and lepton numbers"* [arXiv:1512.06696]. L.M. Carpenter, R. Colburn, J. Goodman, *"Supersoft SUSY Models and the 750 GeV Diphoton Excess, Beyond Effective Operators"* [arXiv:1512.06107]. R. Ding, L. Huang, T. Li, B. Zhu, *"Interpreting $750\,\mathrm{GeV}$ Diphoton Excess with R-parity Violation Supersymmetry"* [arXiv:1512.06560]. F. Wang, L. Wu, J.M. Yang, M. Zhang, *"750 GeV Diphoton Resonance, 125 GeV Higgs and Muon g-2 Anomaly in Deflected Anomaly Mediation SUSY Breaking Scenario"* [arXiv:1512.06715]. B.C. Allanach, P.S.B. Dev, S.A. Renner, K. Sakurai, *"Di-photon Excess Explained by a Resonant Sneutrino in R-parity Violating Supersymmetry"* [arXiv:1512.07645]. J.A. Casas, J.R. Espinosa, J.M. Moreno, *"The 750 GeV Diphoton Excess as a First Light on Supersymmetry Breaking"* [arXiv:1512.07895]. L.J. Hall, K. Harigaya, Y. Nomura, *"$750\,\mathrm{GeV}$ Diphotons: Implications for Supersymmetric Unification"*, JHEP 1603 (2016) 017 [arXiv:1512.07904]. Y. Jiang, Y-Y. Li, T. Liu, *"$750\,\mathrm{GeV}$ Resonance in the Gauged $U(1)'$-Extended MSSM"* [arXiv:1512.09127]. W. Chao, *"The Diphoton Excess from an Exceptional Supersymmetric Standard Model"* [arXiv:1601.00633]. B. Dutta, Y. Gao, T. Ghosh, I. Gogoladze, T. Li, Q. Shafi, J.W. Walker, *"Diphoton Excess in Consistent Supersymmetric SU(5) Models with Vector-like Particles"* [arXiv:1601.00866]. S.F. King, R. Nevzorov, *"$750\,\mathrm{GeV}$ Diphoton Resonance from Singlets in an Exceptional Supersymmetric Standard Model"*, JHEP 1603 (2016) 139 [arXiv:1601.07242]. G. Lazarides, Q. Shafi, *"Diphoton Resonances in $\mathrm{U}(1)_{B-L}$ Extension of MSSM"* [arXiv:1602.07866]. R. Ding, Y. Fan, L. Huang, C. Li, T. Li, S. Raza, B. Zhu, *"Systematic Study of Diphoton Resonance at 750 GeV from Sgoldstino"* [arXiv:1602.00977]. D. Bardhan, P. Byakti, D. Ghosh, T. Sharma, *"The $750\,\mathrm{GeV}$ diphoton resonance as an sgoldstino: a reappraisal"* [arXiv:1603.05251]. P. Baratella, J. Elias-Miro, J. Penedo, A. Romanino, *"A closer look to the sgoldstino interpretation of the diphoton excess"* [arXiv:1603.05682]. H.P. Nilles, M.W. Winkler, *"750 GeV Diphotons and Supersymmetric Grand Unification"* [arXiv:1604.03598]. D. Choudhury, K. Ghosh, *"The LHC Diphoton excess at $750\,\mathrm{GeV}$ in the framework of the Constrained Minimal Supersymmetric Standard Model"* [arXiv:1605.00013]. A. Djouadi, A. Pilaftsis, *"The $750\,\mathrm{GeV}$ Diphoton Resonance in the MSSM"* [arXiv:1605.01040].

[11] E. Molinaro, F. Sannino, N. Vignaroli, *"Minimal Composite Dynamics versus Axion Origin of the Diphoton excess"* [arXiv:1512.05334]. J.M. No, V. Sanz, J. Setford, *"See-Saw Composite Higgses at the LHC: Linking Naturalness to the 750 GeV Di-Photon Resonance"* [arXiv:1512.05700]. A. Belyaev, G. Cacciapaglia, H. Cai, T. Flacke, A. Parolini, H. Serdio, *"Singlets in Composite Higgs Models in light of the LHC di-photon searches"* [arXiv:1512.07242].

[12] O. Antipin, M. Mojaza, F. Sannino, *"A natural Coleman-Weinberg theory explains the diphoton excess"* [arXiv:1512.06708].

[13] L. Marzola, A. Racioppi, M. Raidal, F.R. Urban, H. Veermae, *"Non-minimal CW inflation, electroweak symmetry breaking and the 750 GeV anomaly"*, JHEP 1603 (2016) 190 [arXiv:1512.09136].

[14] G.M. Pelaggi, A. Strumia, E. Vigiani, *"Trinification can explain the di-photon and di-boson LHC anomalies"*, JHEP 1603 (2016) 025 [arXiv:1512.07225].

[15] K. Kannike, *"Vacuum Stability Conditions From Copositivity Criteria"*, Eur. Phys. J. C72 (2012) 2093 [arXiv:1205.3781].

[16] R. Franceschini, G.F. Giudice, J.F. Kamenik, M. McCullough, F. Riva, A. Strumia, R. Torre, *"Digamma, what next?"* [arXiv:1604.06446].

[17] ATLAS Collaboration, *"Search for squarks and gluinos in final states with jets and missing transverse momentum at* $\sqrt{s} = 13\,\mathrm{TeV}$ *with the ATLAS detector"* [arXiv:1605.03814].

[18] J. R. Ellis, M. K. Gaillard and D. V. Nanopoulos, *"A Phenomenological Profile of the Higgs Boson"*, Nucl. Phys. B 106 (1976) 292. M. A. Shifman, A. I. Vainshtein, M. B. Voloshin and V. I. Zakharov, *"Low-Energy Theorems for Higgs Boson Couplings to Photons"*, Sov. J. Nucl. Phys. 30 (1979) 711 [Yad. Fiz. 30 (1979) 1368].

[19] D. Buttazzo, G. Degrassi, P. P. Giardino, G. F. Giudice, F. Sala, A. Salvio, A. Strumia, *"Investigating the near-criticality of the Higgs boson"*, JHEP 1312 (2013) 089 [arXiv:1307.3536].

[20] G. Degrassi, S. Di Vita, J. Elias-Miro, J. R. Espinosa, G. F. Giudice, G. Isidori, A. Strumia, *"Higgs mass and vacuum stability in the Standard Model at NNLO"*, JHEP 1208 (2012) 098 [arXiv:1205.6497].

[21] V. Branchina, E. Messina, *"Stability, Higgs Boson Mass and New Physics"*, Phys. Rev. Lett. 111 (2013) 241801 [arXiv:1307.5193].

[22] M. Dhuria, G. Goswami, *"Perturbativity, vacuum stability and inflation in the light of 750 GeV diphoton excess"* [arXiv:1512.06782]. J. Zhang, S. Zhou, *"Electroweak Vacuum Stability and Diphoton Excess at 750 GeV"*, Chin. Phys. C 40 (2013) 081001 [arXiv:1512.07889]. A. Salvio, A. Mazumdar, *"Higgs Stability and the 750 GeV Diphoton Excess"*, Phys. Lett. B755 (2016) 469 [arXiv:1512.08184]. Y. Hamada, T. Noumi, S. Sun, G. Shiu, *"An O(750) GeV Resonance and Inflation"* [arXiv:1512.08984]. S. F. Ge, H. J. He, J. Ren, Z. Z. Xianyu, *"Realizing Dark Matter and Higgs Inflation in Light of LHC Diphoton Excess"*, Phys. Lett. B757 (2016) 480 [arXiv:1602.01801]. K. J. Bae, M. Endo, K. Hamaguchi, T. Moroi, *"Diphoton Excess and Running Couplings"*, Phys. Lett. B757 (2016) 493 [arXiv:1602.03653].

[23] A. Salvio, F. Staub, A. Strumia, A. Urbano, *"On the maximal diphoton width"*, JHEP 1603 (2016) 214 [arXiv:1602.01460].

[24] H. Ishida, S. Matsuzaki, Y. Yamaguchi, *"Invisible Axion-Like Dark Matter from Electroweak Bosonic Seesaw"* [arXiv:1604.07712].

[25] K. Harigaya, Y. Nomura, *"Composite Models for the* $750\,\mathrm{GeV}$ *Diphoton Excess"*, Phys. Lett. B754 (2016) 151 [arXiv:1512.04850]. Y. Nakai, R. Sato, K. Tobioka, *"Footprints of New Strong Dynamics via Anomaly and the* $750\,\mathrm{GeV}$ *Diphoton"*, Phys. Rev. Lett. 116 (2016) 151802

[arXiv:1512.04924]. A. Pilaftsis, *"Di-photon Signatures from Heavy Axion Decays at the CERN Large Hadron Collider"*, Phys. Rev. D93 (2016) 015017 [arXiv:1512.04931]. Y. Bai, J. Berger, R. Lu, *"750GeV dark pion: Cousin of a dark G -parity odd WIMP"*, Phys. Rev. D93 (2016) 076009 [arXiv:1512.05779]. A. Belyaev, G. Cacciapaglia, H. Cai, T. Flacke, A. Parolini, H. Serdio, *"Singlets in Composite Higgs Models in light of the LHC di-photon searches"* [arXiv:1512.07242]. L. Bian, N. Chen, D. Liu, J. Shu, *"Hidden confining world on the* $750\,\mathrm{GeV}$ *diphoton excess"*, Phys. Rev. D93 (2016) 095011 [arXiv:1512.05759]. E. Molinaro, F. Sannino, N. Vignaroli, *"Minimal Composite Dynamics versus Axion Origin of the Diphoton excess"* [arXiv:1512.05334]. E. Molinaro, F. Sannino, N. Vignaroli, *"Minimal Composite Dynamics versus Axion Origin of the Diphoton excess"* [arXiv:1512.05334] K. Harigaya, Y. Nomura, *"A Composite Model for the* $750\,\mathrm{GeV}$ *di-photon Excess"* [arXiv:1602.01092]. M. Redi, A. Strumia, A. Tesi, E. Vigiani, *"Di-photon resonance and Dark Matter as heavy pions"*, JHEP 1605 (2016) 078 [arXiv:1602.07297]. K. Harigaya, Y. Nomura, *"Hidden Pion Varieties in Composite Models for Diphoton Resonances"* [arXiv:1603.05774]. Y. Bai, V. Barger, J. Berger, *"Color-octet Companions of a* $750\,\mathrm{GeV}$ *Heavy Pion"* [arXiv:1604.07835].

[26] O. Antipin, M. Redi, A. Strumia, E. Vigiani, *"Accidental Composite Dark Matter"*, JHEP 1507 (2015) 039 [arXiv:1503.08749].

[27] M. Redi, A. Strumia, A. Tesi, E. Vigiani, *"Di-photon resonance and Dark Matter as heavy pions"* [arXiv:1602.07297].

[28] F. Feruglio, B. Gavela, K. Kanshin, P.A.N. Machado, S. Rigolin, S. Saa, *"The minimal linear sigma model for the Goldstone Higgs"* [arXiv:1603.05668].

[29] E. Megias, O. Pujolas, M. Quiros, *"On dilatons and the LHC diphoton excess"* [arXiv:1512.06106]. J. Cao, L. Shang, W. Su, Y. Zhang, J. Zhu, *"Interpreting the* $750\,\mathrm{GeV}$ *diphoton excess in the Minimal Dilaton Model"* [arXiv:1601.02570]. B. Agarwal, J. Isaacson, K.A. Mohan, *"Minimal Dilaton Model and the Diphoton Excess"* [arXiv:1604.05328].

[30] R. Jinno, M. Takimoto, *"Probing classically conformal* $B - L$ *model with gravitational waves"* [arXiv:1604.05035].

[31] G. Servant, *"Baryogenesis from Strong* CP *Violation and the QCD Axion"*, Phys. Rev. Lett. 113 (2014) 171803 [arXiv:1407.0030].